Chapter 01: Gametogenesis

Introduction

The human reproductive system is the only biological system that remains inactive until puberty. This unique feature ensures that the body matures before engaging in reproductive activities. The gonads, which include the **testes in males** and the **ovaries in females**, serve two primary functions: the production of gametes and the secretion of sex hormones. Gametes are specialized reproductive cells responsible for transmitting genetic information to the next generation. Additionally, sex hormones play a crucial role in the regulation of reproductive functions and the development of secondary sexual characteristics.

Gametogenesis

Gametogenesis is the biological process responsible for the formation of gametes, ensuring genetic continuity across generations. This process involves cell division and differentiation to produce **haploid (n) gametes** from **diploid (2n) precursor cells**. In males, the process is called **spermatogenesis**, while in females, it is known as **oogenesis**. Spermatogenesis takes place in the **seminiferous tubules** of the testes and begins at puberty, continuing throughout a man's lifetime. In contrast, oogenesis occurs in the ovaries and is largely completed before birth, with the maturation of ova occurring cyclically from puberty until menopause.

Dr. OUARABI.L

Part 01: Spermatogenesis

I. Male Reproductive System

The male reproductive system is a complex network of organs that work together to produce, store, and transport spermatozoa. It consists of:

- **Two testes** – The primary male reproductive organs responsible for sperm production and hormone secretion.
- **Excretory genital ducts** – Includes the **epididymis**, **vas deferens**, and **ejaculatory ducts**, which facilitate sperm transport.
- **Accessory glands** – The **seminal vesicles**, **prostate gland**, and **bulbourethral glands**, which produce seminal fluid to nourish and protect sperm.
- **Penis** – The organ responsible for delivering sperm into the female reproductive tract.




Figure 4. Male reproductive system anatomy

1. The Testes

The **testes** are oval-shaped structures located within the **scrotum**, an external pouch that regulates their temperature. They measure approximately **4 cm in length** and weigh between **10 to 15g** each. Their placement outside the abdominal cavity provides a cooler environment necessary for efficient sperm production.




Figure 5. Testis anatomy

The **testis perform two essential functions**:

- **Exocrine function** – The production of male gametes (**spermatogenesis**).
- **Endocrine function** – The secretion of androgenic hormones, primarily **testosterone**, which is crucial for male reproductive health.


Each testis is encased in a dense fibrous layer called the **tunica albuginea**, which divides the testis into **200-300 lobules**. Each lobule contains **1 to 3 seminiferous tubules**, the site of sperm production. The testicular structure also includes interstitial cells responsible for hormone secretion.

2. The Seminiferous Tubules

Seminiferous tubules are highly coiled structures that serve as the site of sperm cell formation. Each tubule measures between **30 to 70 cm in length** and **200 µm in diameter**. Within these tubules, two main types of cells can be found:

- **Germ cells** – Present at different stages of spermatogenesis, responsible for producing spermatozoa.
- **Sertoli cells** – Somatic cells that provide structural and metabolic support to developing germ cells.

Between the seminiferous tubules lies the **interstitial tissue**, which houses **Leydig cells**. These specialized cells secrete **testosterone**, the primary male sex hormone necessary for spermatogenesis and the development of secondary sexual characteristics.



Dr OUARABI.L

Figure 6. Cross-section of a seminiferous tube

II. Spermatogenesis

1. Definition

Spermatogenesis is the **continuous and highly regulated process** by which sperm cells are produced from precursor cells within the testes. It begins at **puberty** and persists throughout a man's lifetime. The entire process takes approximately **74 days** to complete and involves several stages of **cell division (mitosis and meiosis)**, followed by **morphological differentiation (spermiogenesis)** to generate functionally mature spermatozoa.

2. Pathway of Sperm Cells

The development of sperm occurs within the **seminiferous tubules** and follows a distinct pathway. As cells progress through the stages of spermatogenesis, they move closer to the lumen of the tubule. Once fully matured, sperm cells are released into the lumen and travel toward the **epididymis**, where they undergo final maturation and are stored until ejaculation.

3. Stages of Spermatogenesis

Stage 1: Multiplication (Mitosis)

- **Spermatogonia** are diploid stem cells that serve as the foundation of spermatogenesis.
- They divide by **mitosis**, ensuring a continuous supply of germ cells.
- Some of the resulting cells remain as **stem cells**, while others differentiate into **primary spermatocytes**.

Stage 2: Growth

- **Primary spermatocytes** increase in size as they accumulate cytoplasmic content and organelles.
- These enlarged cells are referred to as **auxocytes**.

Stage 3: Maturation (Meiosis)

- **Meiosis I**: Each **primary spermatocyte** divides into two **secondary spermatocytes**.
- **Meiosis II**: Each secondary spermatocyte undergoes another division, yielding **haploid spermatids**.




Figure 7. Steps of spermatogenesis

Dr OUARABI.L


Stage 4: Differentiation (Spermiogenesis)

- Spermatids undergo structural modifications to become spermatozoa:
  - The **Golgi apparatus** forms the **acrosome**, which contains enzymes to penetrate the egg.
  - The **centriole** elongates to form the **flagellum**, enabling sperm motility.
  - **Mitochondria** cluster in the midpiece, providing the energy required for movement.
  - The cytoplasm is reduced, ensuring a streamlined shape for efficient motility.

4. Structure of Sperm Cells

A fully developed sperm cell comprises:

- **Head** – Contains the **nucleus** with genetic material, and is covered by the **acrosome**, which facilitates egg penetration.
- **Midpiece** – Rich in mitochondria to generate energy for movement.
- **Tail (flagellum)** – Ensures motility and propels the sperm toward the egg.



After **spermiogenesis**, mature spermatozoa enter the **lumen of the seminiferous tubules** in a process called **spermiation**. They are then transported to the **epididymis**, where they gain motility and the ability to fertilize an egg.

Figure 8. Sperm structure




Figure 9. Meiosis on spermatogenesis

III. Hormonal Regulation in Males

The regulation of spermatogenesis is controlled by the **hypothalamic-pituitary-gonadal axis**:

- The **hypothalamus** releases **Gonadotropin-Releasing Hormone (GnRH)**.
- **GnRH** stimulates the pituitary gland to secrete:
  - **FSH (Follicle-Stimulating Hormone)** – Activates Sertoli cells to nourish developing sperm cells.
  - **LH (Luteinizing Hormone)** – Stimulates Leydig cells to produce testosterone.
- **Testosterone** triggers spermatogenesis and promotes secondary sexual traits.




Figure 10. Male hormonal regulation

2. Negative Feedback Mechanism

- **High testosterone levels** inhibit GnRH and pituitary hormone release.
- **Inhibin**, secreted by Sertoli cells, decreases **FSH production**, regulating sperm production.

This feedback system maintains hormonal balance, ensuring optimal sperm production and reproductive health.




Figure 11. Feedback male system

Dr. OUAR