

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/370865204>

Impact of Thermal and Non-Thermal Pasteurization on the Microbial Inactivation of Fruit Juice: Review

Article · May 2023

DOI: 10.35248/2476-2059.23.8.198

CITATIONS

2

12 authors, including:



Ourdia Kernou

Université de Béjaïa

19 PUBLICATIONS 52 CITATIONS

SEE PROFILE



Kenza Bedjaoui

Université de Béjaïa

8 PUBLICATIONS 67 CITATIONS

SEE PROFILE

READS

442



Ghania Kaanin

Université de Béjaïa

7 PUBLICATIONS 29 CITATIONS

SEE PROFILE



Sara Guemouni

Université de Béjaïa

7 PUBLICATIONS 23 CITATIONS

SEE PROFILE

Impact of Thermal and Non-Thermal Pasteurization on the Microbial Inactivation of Fruit Juice: Review

Belbahi Amine¹, Amir Nadir², Kernou Ourdia Nouara^{3*}, Amir Tahi Akila³, Kerdouche Kamelia³, Kaanin Boudraa Ghania³, Bedjaoui Kenza³, Guemouni Sara³, Djerroud Mohellebi Naima³, Debbou Iouknane Nedjima⁴, Boulekbache Lila³, Madani Khodir^{3,5}

¹Department of Microbiology and Biochemistry, University of M' Sila, M' Sila, Algeria; ²Département de Microbiologie, Université de Bejaia, Route de Targa-Ouzemour, Bejaia, 06000, Algeria; ³Laboratoire de Biomathématiques, Biophysique, Biochimie, et Scientométrie (L3BS), Université de Bejaia, Bejaia, 06000, Algeria; ⁴Department of Environment Biological Sciences, University of Bejaia, Route de Targa Ouzemmour, Bejaia, 06000, Algeria; ⁵Centre de recherche en technologie agroalimentaire, Route de targua-ouzemour, Bejaia, Algeria

ABSTRACT

Because of their exceptional qualities and high nutritional content, fruit and vegetable juices are an indispensable component of a healthy diet. The juices must first go through the process of pasteurization, which is the fundamental step in ensuring their quality, safety, and longevity. Although heat pasteurization is the most frequent approach for rendering bacteria inactive, it results in a number of unwanted side effects that contribute to a decline in the juice's overall quality. As a result, a number of non-thermal techniques of pasteurization have been devised, each of which assures the juice's safety while causing very minimal modifications to the juice's qualities. Yet, the high cost of installation and operation presented by these systems is the primary barrier that prevents their widespread use in industrial settings. Methods of non-thermal pasteurization, such as membrane filtration, pulsed electric field, ultraviolet, and sonication treatments, are discussed in this article. Other non-thermal pasteurization techniques are also discussed.

Keywords: Fruit juice; Pasteurization; Membrane filtration; Ultraviolet exposure; Pulsed electric field; Sonication treatment; Microbial inactivation

INTRODUCTION

Fruit juices are the most popular drinks and make up a big part of the food industry's market. This is because they have a unique mix of physical and chemical properties that make them natural and healthy [1]. These low-calorie foods, which are packed with nutrients and bioactive substances like proteins, carbohydrates, polyphenols, enzymes, minerals, fibers, and antioxidants, vitamins, can fit into today's hectic lifestyle [2]. Most acidic fruits are perishable because their high water activity promotes microbial and metabolic activities that ruin them. In addition to sanitary procedures, several treatments are applied during processing and/or storage to prevent the growth of undesirable

bacteria. Fruit juices come in a wide range and are a vital source of complex nutrients for good health. Both foodborne pathogens and microorganisms that cause spoilage can grow well in the presence of these nutrients. *Salmonella*, *Cryptosporidium*, *Listeria monocytogenes*, and *Escherichia coli* are all relevant pathogens for procedures intended to regulate the efficacy of disinfection treatments, depending on the type of juice [3]. Thermal pasteurization and the natural acidity of fruit juices are used in treatment methods. The final product's overall quality might be impacted by unfavorable biochemical and nutritional changes brought on by the heat application. To maximize bacterial inactivation and minimize nutrient loss during thermal pasteurization, there have been several attempts to alter

Correspondence to: Kernou Ourdia Nouara, Laboratoire de Biomathématiques, Biophysique, Biochimie, et Scientométrie (L3BS), Université de Bejaia, Bejaia, 06000, Algeria, E-mail: ourdia.kernou@univ-bejaia.dz

Received: 04-Apr-2023, Manuscript No. JFMSH-23-23074; **Editor assigned:** 06-Apr-2023, Pre QC No. JFMSH-23-23074 (PQ); **Reviewed:** 20-Apr-2023, QC No. JFMSH-23-23074; **Revised:** 27-Apr-2023, Manuscript No. JFMSH-23-23074 (R); **Published:** 04-May-2023, DOI: 10.35248/2476-2059.23.8.198.

Citation: Amine B, Nadir A, Nouara KO, Akila AT, Kamelia K, Ghania KB, et al (2023) Impact of Thermal and Non-Thermal Pasteurization on the Microbial Inactivation of Fruit Juice: Review. J Food Microbial Saf Hyg. 8:198.

Copyright: © 2023 Amine B, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

parameters, including processing time and temperature [4,5]. As a result of the variants' unsuccessful validation, the majority of these methods cannot be applied in the workplace [6].

In order to pasteurize food, a number of non-thermal techniques have been developed, including Ultra Violet (UV) exposure [7-9], high-voltage Pulsed Electric Fields (PEF) [10-12], membrane filtration [13,14], high hydrostatic pressure [15,16], and sonication [17-19]. These technologies have the benefit of preserving food's "fresh-like" qualities while also saving time and energy. The processing of fruit juices, particularly apple juice, has made extensive use of membrane filtration techniques [20]. PEF, which has already been used in an industrial setting, is among the most innovative non-thermal technologies [21]. When treating liquid food, it can inactivate microorganisms [22] and enzymes [23] with a relatively small temperature increase, resulting in a minimal loss of nutrients and minor quality changes. Another non-thermal pasteurization method that is frequently used in food preservation is UV exposure. The DNA of microorganisms can absorb short-wave UV light, which prevents cells from replicating [24].

LITERATURE REVIEW

This article reviews both thermal and non-thermal pasteurization methods for microbial inactivation, including PEF, membrane filtration, UV exposure, and sonication.

Pasteurisation technique in fruit juice preservation

Thermal pasteurization: The most popular method for pasteurizing fruit juice is thermal processing. The most resilient microorganisms of public health concern are reduced by five logs during the pasteurization of juice [25]. This approach depends on the heat that is created outside a meal and then transported into the food *via* conduction and convection processes [26]. By an appropriate time/temperature combination, thermal

pasteurization aims to kill pathogens and significantly lower the amount of spoiling microorganisms [27]. Conventional thermal pasteurization may be categorized into High-Temperature/Short-Time (HTST) and Low Temperature/Long-Time (LTLT) procedures. Whereas fruit juices are subjected to HTST pasteurization, which is done at temperatures of about 72°C with holding periods of 15 s and higher, LTLT pasteurization entails heating a meal at around 63°C for no less than 30 min. Both procedures may damage the taste, color, flavor, and nutritional content of meals [28].

Table 1 provides an overview of current research on the impact of heat pasteurization on the microbiological inactivation of fruit juices. Several fruit liquids were subjected to HTST pasteurization at temperatures ranging from 72°C to 108°C, with treatments lasting always one minute or less. Higher temperatures were more effective in preventing bacterial development, but they also caused a greater drop in the amount of phenolic chemicals in the juice. Other elements, such as the product complexity and microorganisms, may also have an impact on the effectiveness of HTST therapy. In less complicated single juices, HTST pasteurization performs much better than on more intricate/viscous juice products. For instance, a 15-second HTST treatment at 72°C led to a 6.0 log reduction in the number of native microorganisms in apple juice [29]; but only a 3.5 log reduction in fruit smoothies. As compared to the product's mixed natural populations of microorganisms, single species of microorganisms were more sensitive to heat treatment. For instance, a 15-second HTST treatment at 72°C resulted in a 6.3 log decrease in *Escherichia coli* K12 in fruit smoothies [30], but only a 3.5 log reduction in native microorganisms. Technology advancements made it possible to optimize heat processing for greatest effectiveness against microbiological contaminants and the least amount of food quality degradation [31]. Other technologies that aren't thermal, however, provide us with additional options.

Type of juice	Conditions	Microbial inactivation (log ₁₀)	References
Longan	100°C, 1 min	7.0	[32]
Apple	72°C, 26 s	6.0	[29]
Apple	94°C, 26 s	6.7	[33]
Orange	90°C, 1 min	2.5	[34]
Strawberry	90°C, 30 s	N/A	[35]
	90°C, 60 s	N/A	
Apple/cranberry	72°C, 26 s	N/A	[36]
Tomato	98°C, 40 s	N/A	[37]
	108°C, 40 s	N/A	
	128°C, 40 s	N/A	

Fruit smoothie	72°C, 15 s	3.5	[22]
	72°C, 15 s	<i>E. coli</i> 6.3	

Note: N/A, not reported.

Table 1: Effect of thermal pasteurization on microbial inactivation of fruit juice.

Non-thermal pasteurization technologies: For the last 30 years, non-thermal technologies have undergone extensive research [32].

Membrane filtration: The two most common membrane filtration methods for processing fruit juice are Ultra Filtration (UF) and Micro Filtration (MF). They have been used on a commercial scale to clarify fruit liquids. Essentially, the membranes enable tiny molecules like vitamins, salts, carbohydrates, and water to pass through them while retaining big molecules like microorganisms, proteins, lipids, and colloids (UF only). As a result, "cold pasteurized" goods (>5 log decrease or eradication of microorganisms) may be created using this procedure with superior flavors than products that have undergone heat treatment [38, 39].

"Treat and trans" is a term used to describe trans people, depending on the circumstances. The findings showed that while pH, soluble solids and acid content did not change, the smaller pore membrane treatment reduced their variability. Both membranes showed relative color changes, although the bigger pore membrane treatment was easier to see [40].

Techniques for membrane filtration offer numerous benefits over heat processing. They meet the customer's desire for wholesome, natural foods with a long shelf life and no preservatives [39]. Comparing membrane filtering to traditional heat processing, energy savings are also significant. Moreover, it protects the environment by reducing trash production [41]. These benefits have led to the widespread usage of membrane filtering systems in the production of fruit juice. Membrane fouling (rapid decrease of permeate flow) and cleaning are the primary drawbacks of membrane filtration systems, which ultimately result in high expenses for membrane replacement and make the filtration system relatively costly in terms of operating costs and maintenance [42].

Pulsed electric field: PEF pasteurization is a method based on the transmission of millisecond-long pulses of high electric field strength (5-55 kV cm⁻¹) to food [31]. It may be used to produce safe, shelf-stable goods like fruit juice that have great nutritional contents and "fresh-like" qualities. It has been shown that PEF, with substantially less phenolic degradation, may achieve an equal level of microbial inactivation effectiveness as heat pasteurization. Table 2 provides examples of how PEF pasteurization affects the microbial inactivation of fruit juice.

PEF is a method of continuous processing used to inactivate vegetative organisms like *E. coli* and yeast. For the inactivation of bacterial spores, it is ineffective enough, albeit [31]. Although studies on outbreaks have linked fruit juices to the presence of

harmful microorganisms such as *E. coli* O157: H7 [43], PEF could be considered an appropriate treatment strategy for fruit juice preservation. The mechanism of membrane structural or functional damage in microorganism inactivation by high-voltage PEF is similar [31]. When PEF is applied, the buildup of molecules with opposing charges surrounding the membrane may raise the transmembrane potential of cells. The cell membrane becomes thinner when the transmembrane potential is surpassed, leading to reversible or irreversible electroporation [44].

Electric field intensity, treatment duration, temperature, and product composition are just a few of the variables that might determine how effective PEF therapy is at inactivating microorganisms.

Electric field strength and microbial inactivation effectiveness were shown to be significantly correlated [45-47]. With increasing temperature, *Escherichia coli* inactivation in apple juice rose from 2 to 3.3 log. The range of the electric field from 16 kV to 20 kV at 35°C [23]. Higher electric field application, nevertheless, may potentially have negative impacts on food quality [48].

Microbial inactivation is influenced by treatment duration as well. As the number of pulses delivered and the pulse width determine the treatment duration, increasing the number of pulses lengthens the treatment period and enhances microbial inactivation. Also, it was hypothesized that the first 10-20 pulses were when the therapy was most effective [47].

Inactivation of microorganisms and phenolic degradation are both significantly influenced by treatment temperature. PEF application at a cool temperature has been recommended as a way to increase the preservation method's potency. When the inlet temperature increases from 10°C to 50°C, microorganism inactivation rates will rise [45]. The phenolics in apple juice, however, may be thermally degraded at higher temperatures. PEF works better on juices with straightforward compositions than on more intricate multi-juice products. For instance, adding more and more carrot juice to orange juice decreased the PEF's ability to kill *E. coli*, which was likely due to the multicomponent product's more complex matrix [49]. Because of their different cell structures, it was considered that Gram-positive bacteria were usually more resistant to PEF than Gram-negative bacteria [50]. For instance, a tropical fruit smoothie with PEF treatment at 34 kV/cm for 150 s resulted in a 5.4-log reduction of *E. coli* [30]. However, despite orange juice being a less viscous and complex food system, *Staphylococcus aureus* still needed to be treated with PEF under harsher circumstances (40 kV cm⁻¹ for 150 s) to be reduced by 5.5 logs [51].

Type of juice	Conditions	Microbial inactivation (log ₁₀)	References
Longan	32 kV cm ⁻¹ , 90s	2	[32]
Orange	35 kV cm ⁻¹ , 1000 µs	25	[34]
Strawberry	35 kV cm ⁻¹ , 1700 µs	N/A	[35]
Apple	40 kV cm ⁻¹ , 100 µs	5.4	[29]
Apple	35 kV cm ⁻¹ , 4800 µs	<i>L. brevis</i> 6.3	[52]
Apple	35 kV cm ⁻¹ , 4800 µs	<i>S. cerevisiae</i> 4.2	[52]
Water melon	35 kV cm ⁻¹ , 50 µs	N/A	[53]
Fruit smoothie	34 kV cm ⁻¹ , 150 µs	<i>E.coli</i> 5.4	[54]
Orange juice	40 kV cm ⁻¹ , 150 µs	<i>S. aureus</i> 5.5	[22]

Note: N/A, not reported.

Table 2: Examples of fruit juice microbial elimination using PEF pasteurization.

Ultraviolet exposure: Broadly acting as an antimicrobial, UV-C light effectively renders viruses, vegetative bacteria, bacterial spores, conidia, yeasts, and parasites inactive. Many studies have shown the capability of UV-C light to destroy bacteria, viruses, and molds. Some of these studies are included in Table 3.

UV-C radiation's effects on microorganisms, however, varies depending on the species, medium, strain, density, and even the size of the organisms. Additionally, UV-C radiation's ability to prevent bacteria from replicating is due to the dimerization of their thymine bases in DNA strands [55], which is only possible when UV-C radiation can pass through liquid food. Fruit juices absorb 90% of UV-C light within a millimeter of the surface [56]. Researchers have also noted that different UV-C light source-to-product distances, power levels, process times, and product thickness were used to achieve different levels of pathogen inactivation, which had a variety of effects on the juice quality. Highly reactive hydroxyl radicals are produced as a result of UV-C light acting as an oxidizing agent [57]. Temperature has a positive impact on this radical production and the cellular components' subsequent response [58]. The four constituent bases of DNA (Deoxyribonucleic Acid) are adenine, guanine, pyrimidines, and purines (thymine and cytosine). A double-stranded helix connects them all. The pyrimidine bases' ability to absorb UV-C radiation allows for a special photochemical reaction that results in the dimerization of nearby pyrimidines (formation of a chemical bond between the pyrimidines). Thymines are the most common nucleotide to dimerize, but cytosine dimers and thymine-cytosine heterodimers can also form. When the cell goes through mitosis, this structural error in the DNA prevents it from replicating [59]. The capsid proteins can be impacted by UV-C light at higher doses (>1000 mJ/cm²). It is believed that nucleic acids and the size/type of the virion work together to affect how sensitive or resistant viruses are to UV-C [60]. The unique fruit juice's absorption coefficient and travel length determine UV-C fluency. Two key

variables might affect these parameters: (1) the properties of the liquid food, such as turbidity, particle size, viscosity, total soluble solids, suspended particles, and in certain cases pH.

These characteristics are crucial in determining how much UV-C light will be absorbed.

Liquid foods differ from water in a variety of optical, physical, and chemical characteristics [61]; in contrast, factor (2) pertains to the UV-C reactor was used to make the juice sensitive to light, which will ultimately affect UV-C light transmittance, momentum transfer, dose delivery, and therefore microbial inactivation [62]. In turn, these elements will affect how much photo-oxidation occurs in the treated juice. In order to produce effective UV-C pasteurization, it is important to know how UV-C affects the relationship between the absorption coefficient of liquid food, particle interference, and soluble and suspended solids.

Guerrero-Beltrén, et al. [63], and Koutchma, et al. [64], where it was discovered that the absorption coefficient and the efficiency of UV-C inactivation are strongly influenced by the product composition, solid contents, color, and overall chemistry of the food product. Different fruit juices with various levels of turbidity, total soluble solids, pH, and viscosity make a big difference in how to successfully treat them with UV-C light. In addition to reducing the UV-C dose through light scattering, suspended solids may also act as a site for bacterial aggregation on particle surfaces [64]. Therefore, it is important to consider the variations and combinations of the juice's physical characteristics, such as turbidity and particle size. Juice composition has a significant impact on its absorption coefficient. Shah, et al. [65], found in her study that juice's absorption coefficient is related to its turbidity and total soluble solids, which also have an impact on the juice's particle size distribution. Unless they have high porosity, the soluble and suspended solids in the juice are small enough to absorb a lot of

light. Porosity, however, may protect microorganisms, allowing them to endure or even be partially harmed by UV-C processing. Although the amount of suspended solids in juice (or juice matter and pulp) determines its turbidity, this factor also has a significant impact on the juice's viscosity characteristics. The

juice's increased flowability or viscosity will subsequently enhance its rate of flow within the UV-C reactor. Increased flow rates led to more effective mixing and irradiation of the juice. Juice turbidity is negatively correlated with UV-C inactivation efficiency, hence lowering it will increase inactivation rate.

Juices	Microflora	UV-C Dosage	Log Reduction	References
Mango nectar	Aerobic plate count	45 J cm ²	2.7	[63]
Pineapple juice	Aerobic plate count	10.76 mJ cm ²	1.9	[66]
	Yeast and mold	10.76 mJ cm ²	1.4	
	<i>S. Typhimurium</i>	0.000154 L s ⁻¹	3	[67]
	Aerobic plate count	1607 J L ⁻¹	<1.0	[33]
	Yeast and mold	1607 J L ⁻¹	<1.0	
Pummela (<i>Citrus grandis</i> L. Osbeck) Juice	<i>S. Typhimurium</i>	15.45-27.63 mJ cm ²	5.23-9.10	[65]
Starfruit juice	Aerobic plate count	NA	Total inactivation	[68]
Lemon -melon juice mix	<i>E. coli</i> K12	0.44-2.86 mJ cm ²	0.06-6 log	[69]
Apple cider	<i>E. coli</i> 0157:H7	4.45, 6.67, 13-34 mJ cm ²	2.85-4.76	[70]
	Aerobic plate count	14 mJ cm ²	1.8	[71]
	Yeast and mold	14 mJ cm ²	1.4	
	<i>E. coli</i> 0157:H7	8.77-35.11 mJ cm ²	>5.0	[72]
	<i>E. coli</i> K12	19.4 mJ cm ²	<2.0	[73]
Apple juice	<i>L. innocua</i>	2.7 J cm ²	4.8-5.8	[36]
	<i>E. coli</i>	7.7 KJ L ⁻¹	6	[74]
	<i>S. cerevisiae</i>	9.6 KJ L ⁻¹	4	
	<i>L. plantarium</i>	3.9 KJ L ⁻¹	>5.0 (total inactivation)	[73]
	<i>A. acidoterrestres</i>	9.6 KJ L ⁻¹	4	
	<i>E. coli</i> K12	24.9 mJ cm ²	>5.0	
Apple and cranberry juice	<i>P. fermentans</i>	5.3 J cm ²	<2.0	[75]
	<i>E. coli</i>	5.3 J cm ²	6	
Grape juice	<i>S. cerevisiae</i>	138 mJ cm ² , 9 min	5	[76]
	Yeasts	280 mJ cm ² , 24 min	3	
	Lactic acid bacteria	280 mJ cm ² , 24 min	04-Mar	
Grape juice (White)	<i>E. coli</i> K12	0.90 mL s ⁻¹	5.2	[77]

	<i>A.acidoterrestris</i>	0.38-1.31 mW cm ⁻²	5.8	[78]
	<i>B.bruxellensis</i>	1377.0 J L ⁻¹	>5.0	[79]
	<i>S.cerevisiae</i>	3672.0 J L ⁻¹	>5.0	
Grape fruit juice	<i>E.coli</i>	19.0 mJ cm ⁻²	5.1	[80]
	<i>S.cerevisiae</i>	14.0 mJ cm ⁻²	6	
Grape juice (Red)	<i>B.bruxellensis</i>	3672.0 J L ⁻¹	2	[79]
	<i>S.cerevisiae</i>	3672.0 J L ⁻¹	>5.0	
	<i>L.plantarum</i>	3672.0 J L ⁻¹	>5.0	
Pomegranate juice	<i>E.coli</i>	62.4 J mL ⁻¹	6.2	[81]
	Aerobic Plate count	62.4 J mL ⁻¹	1.8	
	Yeast and Mold	62.4 J mL ⁻¹	1.5	
Watermelon juice	Aerobic palte count	2.7-37.5 J mL ⁻¹	1.5	[82]
Passion fruit	Aerobic plate count	NA	0.53	[83]
	Yeast and mold	NA	Total inactivation	

Table 3: Effect of UV-C light to destroy microorganisms in fruit juice.

But, according to Müller, et al. [74], they found blood orange juice with the maximum turbidity (9986 NTU), viscosity (2.74 mPa.s), and linear correlation to an absorption coefficient of 194.3 cm⁻¹ had a higher rate of *L.plantarum* inactivation than naturally murky apple juice with an absorption value of 48.4 cm⁻¹. In a separate investigation, Koutchma, et al [84] found that using a coiled UV-C module, the absorption coefficient of juices seems to have a substantial effect on the inactivation of *E. coli* K12. Nevertheless, it was observed that the inactivation had a decrease of less than 1 log₁₀ with only one run through the UV-C reactor at values of the absorption coefficient greater than 48 cm⁻¹. If the passes are increased, the log decrease may be raised, exposing the juice to more UV-C light energy and killing the microorganisms. This observation was supported by *E. coli* K12 inactivation in apple cider (turbidity=1383 NTU, α =57 cm⁻¹), after six reactor runs at a flow rate of 75 L/min, there has been a 1 log₁₀ decrease. UV-C lethality against microorganisms may be enhanced depending on the pass duration and the quantity of UV-C absorbed by the fruit juice.

Combination of PEF and UV: When used as pasteurization techniques, PEF and UV both have drawbacks. Bacterial spores cannot be inactivated using PEF technology, and fruit juices' reduced medium transparency makes UV radiation less effective. On the other hand, employing PEF in conjunction with UV exposure may have synergistic benefits since the microbial inactivation processes of PEF and UV radiation are distinct [29,31,48]. It was found that PEF, followed by UV exposure, had the highest effectiveness in maintaining the quality of blended juice while also reducing microorganisms. But in a different study, Gachovska, et al. [48], no differences in microbial inactivation (5.3 log) between various orders of combined treatment were found.

Table 4 provides a summary of these findings. Overall, phenolic degradation is less affected by hurdle technology than microbial inactivation. According to some reports, PEF followed by UV exposure is preferable to PEF followed by PEF [29,85].

Type juice	Microbial inactivation (log10)					References
	Treatment ^a	PEF	UV	PEF+UV	UV+PEF	
Apple	1	5.4	2.2	6.2	7.1	[29]
Apple	2	58	2.2	N/A	8.3	[54]

Apple	3	4.87	3.46	5.35	5.3	[48]
Apple/cranberry	4	4.5	1.6	N/A	6	[75]
Apple/cranberry	5	N/A	N/A	N/A	N/A	[85]

Note: N/A, not reported. ^aTreatment 1-PEF: 40 Kv cm⁻¹ , 100µs; UV: 254 nm, 30 min, 30 W. Treatment 2-PEF: 36 kV cm⁻¹ , 100 µs; W: 254 nm, 30 W, 30 min. Treatment 3-PEF: 60 Kv min⁻¹ flow rate, 40 µs; UV: 253.7 nm, 2.9 s, 25 W. Treatment 4-PEF: 34 kV cm⁻¹ , 93 µs; UV: 254 nm, 30 s, 35 W. Treatment 5-PEE: 34 kV cm⁻¹ , 93 µs; UV: 254 nm, 30s, 35 W.

Table 4: Effect of PEF and UV pasteurization combined on fruit juice's microbial inactivation.

Sonication Treatment (US): US is one of the non-thermal pasteurization methods used in the processing of liquid foods, particularly juices, purees, and smoothies. In its simplest form, ultrasound is sound pressure waves over 20 kHz [86]. Ultrasonic devices generally run at 20 kHz-10 MHz. Powerful ultrasound, which runs between 20 and 100 kHz, may generate cavitation to

kill germs [87]. Yet, the ultrasonic procedure kills germs through cavitation caused by pressure changes from ultrasonic vibrations.

In article of Kernou, et al. [19], the impact of ultrasound on microorganisms is very well described. Numerous studies have demonstrated that ultrasound can render pathogenic and spoilage microorganisms in fruit juice inactive (Table 5).

Sample	Experimental parameters	Effect of ultrasound	References
Inactivation of <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> in pomegranate juice	20 kHz, amplitude levels of 50, 75 and 100%; 0, 3, 6, 9, 12 and 15 min , 25 ±1°C	15 minutes at 100% amplitude level lowered levels by 3.47 log CFU/mL and 1.86 log CFU/mL, respectively.	[88]
<i>Fusarium oxysporum</i> in orange juice	20kHz, 130 W, amplitude 40-100%, pulse (2 and 10 s), 2-10 min.	Citrus extract and benzoate controlled the growth of <i>F. oxysporum</i> in orange juice and to achieve a decrease of 5 log CFU/mL for at least 14 days.	[89]
<i>Saccharomyces cerevisiae</i> , <i>Pichia membranifigans</i> , <i>Wickerhamomyces anomalus</i> , <i>Aygosaccharomyces bailii</i> , <i>ygosccharomyces rouxii</i> , <i>Candida norvegica</i> in strawberry, orange, apple, pineapple and red-fruits juice.	20 kHz , 130 W amplitude 20-60%, pulse (2 and 6 s), 2-10 min.	Reductions of spoilage organisms	[90, 91]
<i>Escherichia coli</i> in cactus pear juice	20 kHz, 1500W (60%, 70%, 80%, and 90% amplitude), 1, 3 and 5min	In both fruit juices, total inactivation was shown after 5 minutes of ultrasonic treatment at the majority of amplitude values, as measured over a 5-day period.	[92]
Inactivation of <i>Alicyclobacillus acidoterrestris</i> spores in orange juice	24 kHz, 460 W cm ⁻² , 33 W and 105 W cm ⁻² , 162W.	Thermosonication requied at least 8°C lower temperature than thermal treatments to achieve the same spore inactivation. (D75°C- value of 49min for 20.2W/mL vs. 217 min for 0.33W/mL). (D85°C - value decreased from 69 to 29min)	[93]
Inactivation of <i>Alicyclobacillus acidoterrestris</i> spores and <i>Saccharomyces cerevisiae</i> in natural squeezed apple juices	20 kHa, 600 W and 95.2 µm wave amplitude; 10 or 30 min at 20, 30, or 44 1°C and pulsed light (PL) (Xenon 1 and 3 pulses/s; 0.1 m distances ; 2.4-71.6J cm ⁻² ; initial temperature 20, 30, 40 1°C	For <i>S. cerevisiae</i> , there were 3.0 log cycles of spore reduction in commercial apple juice and 2 log cycles in natural juice. Reduction cycles of 6.4 and 5.8 log were obtained.	[94, 95]

<i>Escherichia coli</i> ATCC 35218, <i>Salmonella enteridis</i> MA44 and <i>Saccharomyces cerevisiae</i> KE 162 and indigenous flora in commercial (CAJ) and freshly pressed (NAJ) apple juice	20 kHz, 600 W and 95.2 μm wave amplitude; pulsed light 0.73 J cm^{-2} , 155 mL/min	3.7-6.3 log reductions of inoculates microorganisms. Browning development during storage was prevented.	[95]
The inactivation of <i>Listeria monocytogenes</i> and <i>Escherichia coli</i> suspended in apple and orange juices	35°C, 110 μm , 200 kPa	Increase of 116 W increased the inactivation rate approximately 10-folds in both juices	[96]
<i>Escherichia coli</i> 0157: H7 and <i>Salmonella enteridis</i> in mango juice	25 kHz, 200 W, 50 and 60°C for 10 and 7 min respectively	Inactivation rate was different for both pathogens	[97]
<i>Escherichia coli</i> in apple cider	20 kHz, 100 kPa, 59°C, 4 min	5- log reduction was achieved	[98]
<i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> in pomegranate juice	20 kHz amplitude (50, 75 and 100%) and times : 0, 6, 12, 18, 24 and 30 min	More than a 5 log inactivation of <i>E.coli</i> and 1.36 log inactivation of <i>S.cerevisiae</i>	[99]
<i>Candida parapsilosis</i> and <i>Rhodotorula glutinis</i> in cloudy apple juice	24 kHz, 400 W, amplitude 100%, 35°C, 360 s/100mL.	Shelf life of sonication juices around 21 days with flavour maintained	[100]

Table 5: The impact of ultrasound on sterilization and its parameters.

DISCUSSION

In terms of total soluble solids, pH, and titratable acidity, the survival and development of *Escherichia coli* in cactus pear juice were assessed over a period of five days [92]. After 5 minutes of ultrasonic treatment at 20 kHz, 1500 W, the majority of amplitude levels (60%, 70%, 80%, and 90%) showed complete inactivation. In cloudy apple juice, the microbiological shelf life of the yeasts, *Candida parapsilosis* and *Rhodotorula glutinis* was also investigated [100]. Acoustic waves reduced aerobic mesophilic counts and psychrophilic bacteria in samples by around 3 and 5 log CFU/mL, respectively. Ultrasonic baths may sometimes have little to no impact on the inactivation of bacteria. The survival rate decrease of *E. coli* contained in orange juice on average (1.3 log for 60 min of sonication) is not significantly affected by ultrasonic treatment (42 kHz) [17].

CONCLUSION

Fruit juice has traditionally been preserved by a process known as thermal pasteurization, which has been shown to be highly effective. However, the heat that is applied can result in unwelcome phenolic degradation in addition to other chemical changes in the constituents of the juice. There is hope for the long-term storage of fruit juice in non-thermal methods such as membrane filtration, photoelectric and ultraviolet (UV) irradiation, and sonication.

REFERENCES

1. Kaur C, Kapoor HC. Antioxidants in fruits and vegetables-The millennium's health. *Int J Food Sci. Technol.* 2001;36(7):703-725.

- Sourri P, Tassou CC, Nychas GJ, Panagou EZ. Fruit juice spoilage by *Alicyclobacillus*: Detection and control methods-A comprehensive review. *Foods.* 2022;11(5):747.
- Lima Tribst AA, de Souza Sant'Ana A, de Massaguer PR. Microbiological quality and safety of fruit juices-past, present and future perspectives. *Crit Rev Microbiol.* 2009;35(4):310-339.
- Corradini MG, Peleg M. Demonstration of the Weibull-Log logistic survival model's applicability to non isothermal inactivation of *E. coli* K12 MG1655. *J Food Prot.* 2004;67:2617-2621.
- Aragao GM, Corradini MG, Normand MD, Peleg M. Evaluation of the Weibull and log normal distribution functions as survival models of *Escherichia coli* under isothermal and non isothermal conditions. *Int J Food Microbiol.* 2007;119(3):243-257.
- Zhang M, Oh JK, Cisneros-Zevallos L, Akbulut M. Bactericidal effects of nonthermal low-pressure oxygen plasma on *S. typhimurium* LT2 attached to fresh produce surfaces. *J Food Eng.* 2013;119(3):425-432.
- Chandra RD, Prihastyanti MN, Lukitasari DM. Effects of pH, high pressure processing, and ultraviolet light on carotenoids, chlorophylls, and anthocyanins of fresh fruit and vegetable juices. *EFood.* 2021;2(3):13-124.
- do Prado DB, dos Anjos Szczerepa MM, Capeloto OA, Astrath NG, Dos Santos NC, Previdelli IT, et al. Effect of ultraviolet (UVC) radiation on spores and biofilms of *Alicyclobacillus* spp. in industrialized orange juice. *Int J Food Microbiol.* 2019;305:108238.
- Niu L, Wu Z, Yang L, Wang Y, Xiang Q, Bai Y. Antimicrobial Effect of UVC light-emitting diodes against *Saccharomyces cerevisiae* and their application in orange juice decontamination. *J Food Prot.* 2021;84(1):139-146.
- Korzeniewska E, Sekulska-Nalewajko J, Goławski J, Drózd T, Kiełbasa P. Analysis of changes in fruit tissue after the pulsed electric field treatment using optical coherence tomography. *The Eur Phys J Appl. Phys.* 2020;91(3):30902.
- Salehi F. Physico-chemical properties of fruit and vegetable juices as affected by pulsed electric field: A review. *Int J Food Prop.* 2020;23(1):1036-1050.

12. Dziadek K, Kopeć A, Drózd T, Kielbasa P, Ostafin M, Bulski K, et al. Effect of pulsed electric field treatment on shelf life and nutritional value of apple juice. *J Food Sci. Technol.* 2019;56:1184-1191.
13. Lu C, Bao Y, Huang JY. Fouling in membrane filtration for juice processing. *Curr Opin Food Sci.* 2021;42:76-85.
14. Castro-Muñoz R, Fila V, Barragán-Huerta BE, Yáñez-Fernández J, Piña-Rosas JA, Arboleda-Mejía J. Processing of Xoconostle fruit (*Opuntia joconostle*) juice for improving its commercialization using membrane filtration. *J Food Process. Preserv.* 2018;42(1):e13394.
15. Huang HW, Hsu CP, Wang CY. Healthy expectations of high hydrostatic pressure treatment in food processing industry. *J Food Drug Anal.* 2020;28(1):1-3.
16. Aganovic K, Hertel C, Vogel RF, Johne R, Schlüter O, Schwarzenbolz U, Jäger H, et al. Aspects of high hydrostatic pressure food processing: Perspectives on technology and food safety. *Compr Rev Food Sci Food Saf.* 2021;20(40):3225-3266.
17. Kernou ON, Belbahi A, Amir A, Bedjaoui K, Kerdouche K, Dairi S, et al. Effect of sonication on microwave inactivation of *Escherichia coli* in an orange juice beverage. *J Food Process Eng.* 2021;44(5):e13664.
18. Kernou ON, Azzouz Z, Belbahi A, Kerdouche K, Kaanin-Boudraa G, Amir A, et al. Inactivation of *Escherichia coli* in an orange juice beverage by combined ultrasonic and microwave treatment. *Foods.* 2023;12(3):666.
19. Kernou O-N, Belbahi A, Kaanin-Boudraa G, Adel K, Madani PK. A Review: Ultrasound-microwave technologies as alternative methods for inactivation bacterias in fruit juice. *Int J Anal Chem.* 2022;8(2): 31-40p.
20. Sawitri A, Munir MM, Edikresnha D, Sandi A, Fauzi A, Rajak A, et al. An Investigation on bilayer structures of electrospun polyacrylonitrile nanofibrous membrane and cellulose membrane used as filtration media for apple juice clarification. *Materials Research Express.* 2018;5(5):054003.
21. Zhu Y, Zhang M, Mujumdar AS, Liu Y. Application advantages of new non-thermal technology in juice browning control: A comprehensive review. *Food Rev Int.* 2022;1-22.
22. Walkling-Ribeiro M, Noci F, Cronin DA, Lyng JG, Morgan DJ. Shelf life and sensory attributes of a fruit smoothie-type beverage processed with moderate heat and pulsed electric fields. *LWT-Food Sci Technol.* 2010;43(7):1067-1073.
23. Liu Z, Zhao L, Zhang Q, Huo N, Shi X, Li L, et al. Proteomics-based mechanistic investigation of *Escherichia coli* inactivation by pulsed electric field. *Front Microbiol.* 2019;10:2644.
24. Baysal AH. Short-wave ultraviolet light inactivation of pathogens in fruit juices. *Fruit Juices.* 2018;pp.463-510.
25. Food and Drug Administration. Hazard analysis and critical control point (HACCP); Procedures for the safe and sanitary processing and importing of juice. *Fed. Regist.* 2001;66:6137-6202.
26. Pereira RN, Vicente AA. Environmental impact of novel thermal and non-thermal technologies in food processing. *Food Research International.* 2010;43(7):1936-1943.
27. Ortega-Rivas E. Processing effects on safety and quality of foods. CRC Press. 2009.
28. Charles-Rodríguez AV, Nevárez-Moorillón GV, Zhang Q, Ortega-Rivas EH. Comparison of thermal processing and pulsed electric fields treatment in pasteurization of apple juice. *Food Bioprod Process.* 2007;85(2):93-97.
29. Noci F, Riener J, Walkling-Ribeiro M, Cronin DA, Morgan DJ, Lyng JG. Ultraviolet irradiation and Pulsed Electric Fields (PEF) in a hurdle strategy for the preservation of fresh apple juice. *J Food Eng.* 2008;85(1):141-146.
30. Walkling-Ribeiro M, Noci F, Cronin DA, Lyng JG, Morgan DJ. Inactivation of *Escherichia coli* in a tropical fruit smoothie by a combination of heat and pulsed electric fields. *J Food Sci.* 2008;73(8):M395-M399.
31. Lado BH, Yousef AE. Alternative food-preservation technologies: Efficacy and mechanisms. *Microbes Infect.* 2002;4(4):433-440.
32. Zhang YI, Gao BE, Zhang M, Shi J, Xu Y. Pulsed electric field processing effects on physicochemical properties, flavor compounds and microorganisms of longan juice. *J Food Process. Preserv.* 2010;34(6):1121-1138.
33. Keyser M, Müller IA, Cilliers FP, Nel W, Gouws PA. Ultraviolet radiation as a non-thermal treatment for the inactivation of microorganisms in fruit juice. *Innov Food Sci Emerg Technol.* 2008;9(3): 48-354.
34. Elez-Martinez P, Martin-Belloso O. Effects of high intensity pulsed electric field processing conditions on vitamin C and antioxidant capacity of orange juice and gazpacho, a cold vegetable soup. *Food Chem.* 2007;102(1):201-209.
35. Odriozola-Serrano I, Soliva-Fortuny R, Martín-Belloso O. Phenolic acids, flavonoids, vitamin C and antioxidant capacity of strawberry juices processed by high-intensity pulsed electric fields or heat treatments. *Eur Food Res Technol.* 2008;228:239-248.
36. Caminiti IM, Palgan I, Muñoz A, Noci F, Whyte P, Morgan DJ, et al. The effect of ultraviolet light on microbial inactivation and quality attributes of apple juice. *Food Bioproc Tech.* 2012;5:680-686.
37. Pérez-Conesa D, García-Alonso J, García-Valverde V, Iniesta M-D, Jacob K, Sánchez-Siles LM, et al. Changes in bioactive compounds and antioxidant activity during homogenization and thermal processing of tomato puree. *Innov Food Sci Emerg Technol.* 2009;10(2):179-188.
38. Cassano A, Drioli E, Galaverna G, Marchelli R, Di Silvestro G, Cagnasso P. Clarification and concentration of citrus and carrot juices by integrated membrane processes. *J Food Eng.* 2003;57(2): 153-163.
39. Rektor A, Vatai G. Application of membrane filtration methods for must processing and preservation. *Desalination.* 2004;162:271-277.
40. Zárate-Rodríguez E, Ortega-Rivas E, Barbosa-Cánovas GV. Effect of membrane pore size on quality of ultrafiltered apple juice. *Int. J Food Sci. Technol.* 2001;36(6):663-667.
41. Gergely S, Bekassy-Molnar E, Vatai G. The use of multiobjective optimization to improve wine filtration. *J Food Eng.* 2003;58(4): 311-316.
42. He Y, Ji Z, Li S. Effective clarification of apple juice using membrane filtration without enzyme and pasteurization pretreatment. *Sep Purif Technol.* 2007;57(2):366-373.
43. Moore K, Damrow T. Outbreak of acute gastroenteritis attributable to *Escherichia coli* serotype O104: H21-Helena, Montana. *MMWR Morb Mortal Wkly Rep.* 1995;44(27):501-503.
44. Coster HG, Zimmermann U. The mechanism of electrical breakdown in the membranes of *Valonia utricularis*. *J Membr Biol.* 1975;22:73-90.
45. Huang K, Wang J. Designs of pulsed electric fields treatment chambers for liquid foods pasteurization process: A review. *J Food Eng.* 2009;95(2):227-239.
46. Toepfl S, Heinz V, Knorr D. High intensity pulsed electric fields applied for food preservation. *Chem Eng Process: Process Intensif.* 2007;46(6):537-546.
47. Alkhafaji SR, Farid M. An investigation on pulsed electric fields technology using new treatment chamber design. *Innov Food Sci Emerg Technol.* 2007;8(2):205-212.
48. Gachovska TK, Kumar S, Thippareddi H, Subbiah J, Williams F. Ultraviolet and pulsed electric field treatments have additive effect on inactivation of *E. coli* in apple juice. *J Food Sci.* 2008;73(9):M412-M417.

49. Rodrigo D, Barbosa-Cánovas GV, Martínez A, Rodrigo M. Weibull distribution function based on an empirical mathematical model for inactivation of *Escherichia coli* by pulsed electric fields. *J Food Prot.* 2003;66(6):1007-1012.
50. Hülshöger H, Pötel J, Niemann EG. Electric field effects on bacteria and yeast cells. *Radiat Environ Biophys.* 1983;22:149-162.
51. Walkling-Ribeiro M, Noci F, Riener J, Cronin DA, Lyng JG, Morgan DJ. The impact of thermosonication and pulsed electric fields on *Staphylococcus aureus* inactivation and selected quality parameters in orange juice. *Food Bioproc Tech.* 2009;2:422-430.
52. Aguilar-Rosas SF, Ballinas-Casarrubias ML, Nevarez-Moorillon GV, Martín-Belloso O, Ortega-Rivas E. Thermal and pulsed electric fields pasteurization of apple juice: Effects on physicochemical properties and flavour compounds. *J Food Eng.* 2007;83(1):41-46.
53. Oms-Oliu G, Odriozola-Serrano I, Soliva-Fortuny R, Martín-Belloso O. Effects of high-intensity pulsed electric field processing conditions on lycopene, vitamin C and antioxidant capacity of watermelon juice. *Food Chem.* 2009;115(4):1312-1319.
54. Walkling-Ribeiro M, Noci F, Cronin DA, Riener J, Lyng JG, Morgan DJ. Reduction of *Staphylococcus aureus* and quality changes in apple juice processed by ultraviolet irradiation, pre-heating and pulsed electric fields. *J Food Eng.* 2008;89(3):267-273.
55. Mukhopadhyay S, Ramaswamy R. Application of emerging technologies to control *Salmonella* in foods: A review. *Food Res. Int.* 2012;45(2):666-677.
56. Sizer CE, Balasubramaniam VM. New intervention processes for minimally processed juices. *Food Technol.* 1999;53(10):64-67.
57. Rosenfeldt EJ, Linden KG, Canonica S, von Gunten U. Comparison of the efficiency of OH radical formation during ozonation and the advanced oxidation processes O₃/H₂O₂ and UV/H₂O₂. *Water Res.* 2006;40(20):3695-3704.
58. Yaun BR, Sumner SS, Eifert JD, Marcy JE. Inhibition of pathogens on fresh produce by ultraviolet energy. *Int J Food Microbiol.* 2004;90(1):1-8.
59. Jagger J. Introduction to research in ultraviolet photobiology. 1967.
60. Sommer R, Pribil W, Appelt S, Gehringer P, Eschweiler H, Leth H, et al. Inactivation of bacteriophages in water by means of non-ionizing (UV-253.7 nm) and ionizing (gamma) radiation: A comparative approach. *Water Res.* 2001;35(13):3109-3116.
61. Falguera V, Pagán J, Ibarz A. Effect of UV irradiation on enzymatic activities and physicochemical properties of apple juices from different varieties. *LWT - Food Sci. Technol.* 2011;44(1):115-119.
62. Gómez-López VM, Koutchma T, Linden K. Ultraviolet and pulsed light processing of fluid foods. Novel thermal and non-thermal technologies for fluid foods. 2012;pp.185-223.
63. Guerrero-Beltrán JA, Barbosa-Cánovas GV. Inactivation of *Saccharomyces cerevisiae* and polyphenoloxidase in mango nectar treated with UV light. *J Food Prot.* 2006;69(2):362-368.
64. Koutchma T. Ultraviolet light in food technology: Principles and applications. CRC press. 2019.
65. Shah NN, Shamsuddin R, Rahman RA, Adzahan NM. Furan development In Dean vortex UVC treated Pummelo (*Citrus grandis* L. Osbeck) fruit juice. *Int Food Res J.* 2016;23.
66. Shamsudin R, Adzahan NM, Yee YP, Mansor A. Effect of repetitive ultraviolet irradiation on the physico-chemical properties and microbial stability of pineapple juice. *Innov Food Sci Emerg Technol.* 2014;23:114-120.
67. Mansor A, Shamsudin R, Adzahan NM, Hamidon MN. Efficacy of ultraviolet radiation as non-thermal treatment for the inactivation of *Salmonella typhimurium* TISTR 292 in pineapple fruit juice. *Agric Agric Sci Procedia.* 2014;2:173-180.
68. Bhat R, Ameran SB, Voon HC, Karim A, Tze LM. Quality attributes of starfruit (*Averrhoa carambola* L.) juice treated with ultraviolet radiation. *Food Chem.* 2011;127(2):641-644.
69. Kaya Z, Yıldız S, Ünlütürk S. Effect of UV-C irradiation and heat treatment on the shelf life stability of a lemon-melon juice blend: Multivariate statistical approach. *Innov Food Sci Emerg Technol.* 2015;29:230-239.
70. Orłowska M, Koutchma T, Kostrzyńska M, Tang J, Defelice C. Evaluation of mixing flow conditions to inactivate *Escherichia coli* in opaque liquids using pilot-scale Taylor-Couette UV unit. *J Food Eng.* 2014;120:100-109.
71. Tandon K, Worobo RW, Churey JJ, Padilla-Zakour OI. Storage quality of pasteurized and UV treated apple cider. *J Food Process. Preserv.* 2003;27(1):21-35.
72. Donahue DW, Canitez N, Bushway AA. UV inactivation of *E. Coli* O157: H7 in apple cider: Quality, sensory and shelf-life analysis. *J Food Process. Preserv.* 2004;28(5):368-387.
73. Outchma TK, Arisi BP. Biodosimetry of *Escherichia coli* UV inactivation in model juices with regard to dose distribution in annular UV reactors. *J Food Sci.* 2004;69(1):FEP14-22.
74. Müller A, Stahl MR, Graef V, Franz CM, Huch M. UV-C treatment of juices to inactivate microorganisms using Dean vortex technology. *J Food Eng.* 2011;107(2):268-275.
75. Palgan I, Caminiti IM, Muñoz A, Noci F, Whyte P, et al. Combined effect of selected non-thermal technologies on *Escherichia coli* and *Pichia fermentans* inactivation in an apple and cranberry juice blend and on product shelf life. *Int J Food Microbiol.* 2011;151(1):1-6.
76. Kaya Z, Unluturk S. Processing of clear and turbid grape juice by a continuous flow UV system. *Innov Food Sci Emerg Technol.* 2016;33:282-288.
77. Unluturk S, Atilgan MR. Microbial safety and shelf life of UV-C treated freshly squeezed white grape juice. *J Food Sci.* 2015;80(8):M1831-M1841.
78. Baysal AH, Molva C, Unluturk S. UV-C light inactivation and modeling kinetics of *Alicyclobacillus acidoterrestris* spores in white grape and apple juices. *Int J Food Microbiol.* 2013;166(3):494-498.
79. Fredericks IN, Du Toit M, Krügel M. Efficacy of ultraviolet radiation as an alternative technology to inactivate microorganisms in grape juices and wines. *Food Microbiol.* 2011;28(3):510-517.
80. Geveke DJ, Torres D. Pasteurization of grapefruit juice using a centrifugal ultraviolet light irradiator. *J Food Eng.* 2012;111(2):241-246.
81. Pala ÇU, Toklucu AK. Effect of UV-C light on anthocyanin content and other quality parameters of pomegranate juice. *J Food Compos Anal.* 2011;24(6):790-795.
82. Feng M, Ghafoor K, Seo B, Yang K, Park J. Effects of ultraviolet-C treatment in Teflon®-coil on microbial populations and physico-chemical characteristics of watermelon juice. *Innov Food Sci Emerg Technol.* 2013;19:133-139.
83. Guevara M, Tapia MS, Gómez-López VM. Microbial inactivation and quality of guava and passion fruit nectars treated by UV-C light. *Food Bioproc Tech.* 2012;5:803-807.
84. Koutchma T, Parisi B, Unluturk SK. Evaluation of UV dose in flow-through reactors for fresh apple juice and cider. *Chem. Eng. Commun.* 2006;193(6):715-728.
85. Caminiti IM, Noci F, Muñoz A, Whyte P, Morgan DJ, Cronin DA, et al. Impact of selected combinations of non-thermal processing technologies on the quality of an apple and cranberry juice blend. *Food Chem.* 2011;124(4):1387-1392.
86. Butz P, Tauscher B. Emerging technologies: Chemical aspects. *Food Res. Int.* 2002;35(2-3):279-284.

87. Piyasena P, Mohareb E, McKellar R. Inactivation of microbes using ultrasound: A review. *Int J Food Microbiol.* 2003;87(3):207-216.
88. Alighourchi H, Barzegar M, Sahari MA, Abbasi S. The effects of sonication and gamma irradiation on the inactivation of *Escherichia coli* and *Saccharomyces cerevisiae* in pomegranate juice. *Iran J Microbiol.* 2014;6(1):51.
89. Bevilacqua A, Sinigaglia M, Corbo MR. Ultrasound and antimicrobial compounds: a suitable way to control *Fusarium oxysporum* in juices. *Food Bioproc Tech.* 2013;6:1153-1163.
90. Bevilacqua A, Speranza B, Campaniello D, Sinigaglia M, Corbo MR. Inactivation of spoiling yeasts of fruit juices by pulsed ultrasound. *Food Bioproc Tech.* 2014;7:2189-2197.
91. Bevilacqua A, Campaniello D, Sinigaglia M, Corbo MR. Combination of ultrasound and antimicrobial compounds towards *Pichia* spp. and *Wickerhamomyces anomalus* in pineapple juice. *LWT - Food Sci. Technol.* 2015;64(2):616-622.
92. Cruz-Cansino ND, Reyes-Hernández I, Delgado-Olivares L, Jaramillo-Bustos DP, Ariza-Ortega JA, Ramírez-Moreno E. Effect of ultrasound on survival and growth of *Escherichia coli* in cactus pear juice during storage. *Braz. J. Microbiol.* 2016;47:431-437.
93. Silva FV. High pressure processing pretreatment enhanced the thermosonication inactivation of *Alicyclobacillus acidoterrestris* spores in orange juice. *Food Control.* 2016;62:365-372.
94. Ferrario M, Alzamora SM, Guerrero S. Study of the inactivation of spoilage microorganisms in apple juice by pulsed light and ultrasound. *Food Microbiol.* 2015;46:635-642.
95. Ferrario M, Guerrero S. Effect of a continuous flow-through pulsed light system combined with ultrasound on microbial survivability, color and sensory shelf life of apple juice. *Innov Food Sci Emerg Technol.* 2016;34:214-224.
96. Guzel BH, Arroyo C, Condón S, Pagán R, Bayindirli A, Alpas H. Inactivation of *Listeria monocytogenes* and *Escherichia coli* by ultrasonic waves under pressure at nonlethal (manosonication) and lethal temperatures (manothermosonication) in acidic fruit juices. *Food Bioproc Tech.* 2014;7:1701-1712.
97. Kiang WS, Bhat R, Rosma A, Cheng LH. Effects of thermosonication on the fate of *Escherichia coli* O157: H7 and *Salmonella Enteritidis* in mango juice. *Lett Appl Microbiol.* 2013;56(4):251-257.
98. Lee H, Kim H, Cadwallader KR, Feng H, Martin SE. Sonication in combination with heat and low pressure as an alternative pasteurization treatment-Effect on *Escherichia coli* K12 inactivation and quality of apple cider. *Ultrason. Sonochem.* 2013;20(4):1131-1138.
99. Pala ÇU, Zorba NN, Özcan G. Microbial inactivation and physicochemical properties of ultrasound processed pomegranate juice. *J Food Prot.* 2015;78(3):531-539.
100. Bastianello E, Montemurro F, Fasolato L, Balzan S, Marchesini G, Contiero B, et al. Volatile compounds and microbial development in sonicated cloudy apple juices: Preliminary results. *CYTA J Food .* 2016;14(1):65-73.